

BRIEF REPORT

Food Addiction and Bulimia Nervosa: New Data Based on the Yale Food Addiction Scale 2.0

Sarah-Kristin de Vries¹ & Adrian Meule^{2,3*}

¹Department of Psychology, University of Potsdam, Germany

²Department of Psychology, University of Salzburg, Austria

³Center for Cognitive Neuroscience, University of Salzburg, Austria

Abstract

Previous research on ‘food addiction’ as measured with the *Yale Food Addiction Scale* (YFAS) showed a large overlap between addiction-like eating and bulimia nervosa. Most recently, a revised version of the YFAS has been developed according to the changes made in the diagnostic criteria for substance use disorder in the *Diagnostic and Statistical Manual of Mental Disorders fifth edition*. The current study examined prevalence and correlates of the YFAS2.0 in individuals with bulimia ($n = 115$) and controls ($n = 341$). Ninety-six per cent of participants with bulimia and 14% of controls received a YFAS2.0 diagnosis. A higher number of YFAS2.0 symptoms was associated with lower interoceptive awareness, higher depressiveness, and higher impulsivity in both groups. However, a higher number of YFAS2.0 symptoms was associated with higher body mass and weight suppression in controls only and not in participants with bulimia. The current study is the first to show a large overlap between bulimia and ‘food addiction’ as measured with the YFAS2.0, replicating and extending findings from studies, which used the previous version of the YFAS. Compensatory weight control behaviours in individuals with bulimia likely alleviate the association between addiction-like eating and higher body mass. Thus, the large overlap between bulimia and ‘food addiction’ should be taken into consideration when examining the role of addiction-like eating in weight gain and obesity. Copyright © 2016 John Wiley & Sons, Ltd and Eating Disorders Association.

Received 14 May 2016; Revised 7 July 2016; Accepted 17 July 2016

Keywords

food addiction; Yale Food Addiction Scale; bulimia nervosa; Eating Disorder Diagnostic Scale; impulsivity

*Correspondence

Adrian Meule, PhD, Department of Psychology, University of Salzburg, Hellbrunner Straße 34, 5020 Salzburg, Austria. Tel: +43 662 8044 5106; Fax: +43 662 8044 5126.

Email: adrian.meule@sbg.ac.at

Published online 30 August 2016 in Wiley Online Library (wileyonlinelibrary.com) DOI: 10.1002/erv.2470

Introduction

‘Food addiction’ refers to the idea that certain foods (e.g. highly processed and high-calorie foods) may have an addictive potential and that some forms of overeating may represent an addicted behaviour (Meule, 2015). Although this concept is controversially discussed in the scientific community (Benton, 2010; Rogers & Smit, 2000; Wilson, 2010; Ziauddeen & Fletcher, 2013), it has received increasing interest in recent years (Meule, 2015). This popularity of the food addiction concept can be, in part, attributed to the development of the *Yale Food Addiction Scale* (YFAS), which was the first standardized self-report measure for the assessment of addiction-like eating (Gearhardt, Corbin, & Brownell, 2009). The YFAS consists of 25 items and asks participants about their eating behaviour during the past year. Importantly, items are based on the diagnostic criteria for substance dependence in the *Diagnostic and Statistical Manual of Mental Disorders (DSM) fourth edition* (American Psychiatric Association, 1994), which were translated to refer to food and eating. Likewise, scoring of the scale allows for calculating a symptom count, that is, the number of endorsed ‘food addiction’

symptoms, which can range between zero and seven symptoms. The scale also asks for a clinically significant impairment or distress resulting from individuals’ eating behaviour. When at least three symptoms are met and a clinically significant impairment or distress is present, individuals receive a ‘food addiction’ diagnosis.

In 2013, the fifth edition of the DSM (DSM-5) was released, which includes revised diagnostic criteria for substance use disorder (American Psychiatric Association, 2013). Specifically, four new criteria were added, and diagnostic thresholds were lowered such that the presence of two symptoms (and a clinically significant impairment or distress) suffices to receive a diagnosis of substance use disorder (for a discussion of the four new criteria in relation to food and eating, see Meule & Gearhardt, 2014b). Given these substantial changes in the diagnostic criteria for substance use disorder, the YFAS has been revised recently (Gearhardt, Corbin, & Brownell, 2016). The YFAS2.0 consists of 35 items for measuring 11 ‘food addiction’ symptoms. Besides having more items, the YFAS2.0 differs from the original YFAS in some other aspects as well (e.g. changes in item wordings and response options).

Studies using the previous version of the YFAS showed that a high percentage of individuals with bulimia nervosa receive a YFAS diagnosis (more than 80%; Gearhardt, Boswell, & White, 2014; Granero et al., 2014; Meule, von Rezori, & Blechert, 2014). Thus, bulimia is the diagnostic group with the highest prevalence of YFAS diagnoses (i.e. prevalence is higher than in individuals with anorexia, binge eating disorder, and obese individuals with or without binge eating disorder; Meule & Gearhardt, 2014a). In the initial validation study of the YFAS2.0, two diverse, yet not nationally representative, samples from the USA were investigated and approximately 14–15% of participants received a YFAS2.0 diagnosis (Gearhardt et al., 2016). However, no data in individuals with bulimia have been reported for the YFAS2.0 yet. Due to the addition of symptoms and changes made in item wording, scoring, and diagnostic thresholds, it remains to be tested if using the YFAS2.0 in individuals with bulimia will produce results similar to the previous version.

In the current study, prevalence of YFAS2.0 diagnoses was examined in individuals with bulimia and in controls by using the German version of the YFAS2.0 (Meule, 2016) in an online questionnaire survey. Based on findings with the previous version of the YFAS (Gearhardt et al., 2014; Granero et al., 2014; Meule et al., 2014), it was expected that a high percentage of individuals with bulimia (more than 80%) would receive a YFAS2.0 diagnosis. Based on findings with the English version of the YFAS2.0 (Gearhardt et al., 2016), it was expected that a significantly lower percentage of the controls (approximately 15%) would receive a YFAS2.0 diagnosis.

Moreover, little is known about the correlates of addiction-like eating in individuals with bulimia. For example, while scores on the YFAS and YFAS2.0 have generally been found to be associated with overweight and obesity (Gearhardt et al., 2016; Pursey, Stanwell, Gearhardt, Collins, & Burrows, 2014), it has been suggested that this relationship may be attenuated in individuals with bulimia because of their compensatory weight-control behaviours (Meule, 2012). Therefore, as a secondary aim, correlates of the YFAS2.0 were examined. Based on the findings with the previous version of the YFAS (Granero et al., 2014; Meule & Gearhardt, 2014a) and with the English version of the YFAS2.0 (Gearhardt et al., 2016), it was expected that a higher number of YFAS2.0 symptoms would be associated with higher body mass index (BMI), a lack of interoceptive awareness, higher depressiveness, and higher impulsivity. As weight suppression (i.e. the difference between highest past weight and current weight) has been associated with weight gain and eating pathology (Schaumberg et al., 2016), it was tested if it was also related to higher 'food addiction' symptomatology. Finally, it was explored if relationships of YFAS2.0 symptoms with BMI, weight suppression, interoceptive awareness, depression, and impulsivity were different in the two groups.

Methods

Participants and procedure

The study was approved by the institutional review board at the University of Salzburg. Participants were recruited via online forums targeted at individuals with eating disorders (e.g. Facebook groups for individuals with bulimia) and via student mailing lists

at several German universities. The study was advertised as a study on 'attitudes and eating behaviour', and it was explicated that both individuals with and without eating disorders were invited to participate. Three ×50€ were raffled among participants who completed the entire set of questions. The study's website at www.soscisurvey.de was visited 1590 times. One-thousand and thirty-eight individuals started the survey and, of these, 663 finished the survey. Applying the website's data quality check, which is based on the time spent on each page, led to exclusion of two participants. Of the remaining 661 participants, 100 received an eating disorder diagnosis other than bulimia based on the *Eating Disorder Diagnostic Scale* (EDDS) — DSM-5 version. As there were only few participants in each diagnostic group, however, these participants were excluded from further analyses. Of the remaining 561 participants, 118 received a diagnosis of bulimia nervosa and 443 participants did not receive an eating disorder diagnosis based on the EDDS. As individuals with bulimia all had a BMI >18.5 kg/m² (as this is one criterion for receiving a diagnosis of bulimia according to the EDDS), control participants with a BMI <18.5 kg/m² were excluded. Finally, as there were only three men with a diagnosis of bulimia, men were excluded from further analyses, resulting in a final sample of $n = 115$ women with bulimia and $n = 341$ female controls without eating disorders (Table 1).

Measures

Sociodemographic and anthropometric data

The participants provided their age (years), sex (male/female), height (cm), current weight (kg), and highest weight (kg) at their current height. BMI was calculated as current weight divided by squared height (m). Weight suppression was calculated by subtracting current BMI from highest BMI (highest weight divided by squared height).

Yale Food Addiction Scale 2.0

The 35-item YFAS2.0 (Gearhardt et al., 2016; Meule, 2016) was used for measuring addiction-like eating. A symptom count can be calculated, which can range between 0 and 11 symptoms. Furthermore, a diagnostic score can be calculated ('food addiction' vs no 'food addiction'), and diagnoses can be further specified as *mild*, *moderate*, or *severe*, depending on the number of symptoms present. Internal consistency was $\alpha = .970$ in the current study.

Eating Disorder Diagnostic Scale — DSM-5 version

The 23-item DSM5 version of the EDDS (Meule & Blechert, 2015; Stice, 2014) was used for diagnosing eating disorders (for agreement with interview measures, see Sysko et al., 2015). This self-report scale generates diagnostic scores for the following disorders: anorexia nervosa, bulimia nervosa, binge eating disorder, atypical anorexia nervosa, low-frequency bulimia nervosa, low-frequency binge eating disorder, purging disorder, and night eating syndrome.

Eating Disorder Inventory-2 — interoceptive awareness subscale

The 10-item interoceptive awareness subscale of the Eating Disorder Inventory-2 (Garner, 1991; Thiel et al., 1997) was used

Table 1 Frequencies of YFAS2.0 diagnoses and means of continuous study variables for individuals with bulimia nervosa and controls without eating disorders

| | Bulimia nervosa (<i>n</i> = 115) | Controls (<i>n</i> = 341) | Test statistic | <i>p</i> |
|--|-------------------------------------|-------------------------------------|----------------------|----------|
| Age (years) | <i>M</i> = 26.4 (<i>SD</i> = 8.76) | <i>M</i> = 26.3 (<i>SD</i> = 7.42) | $t_{(454)} = 0.17$ | .866 |
| Body mass index (kg/m ²) | <i>M</i> = 26.1 (<i>SD</i> = 8.00) | <i>M</i> = 23.1 (<i>SD</i> = 5.00) | $t_{(454)} = 4.71$ | <.001 |
| Weight suppression (kg/m ²) | <i>M</i> = 4.14 (<i>SD</i> = 4.02) | <i>M</i> = 2.30 (<i>SD</i> = 2.58) | $t_{(454)} = 5.68$ | <.001 |
| Yale Food Addiction Scale 2.0 (diagnoses) | <i>n</i> = 110 (95.7%) | <i>n</i> = 49 (14.4%) | $\chi^2_{(1)} = 250$ | <.001 |
| Yale Food Addiction Scale 2.0 (symptoms) | <i>M</i> = 8.91 (<i>SD</i> = 2.22) | <i>M</i> = 2.00 (<i>SD</i> = 2.79) | $t_{(454)} = 24.1$ | <.001 |
| Eating Disorder Inventory-2 (interoceptive awareness) | <i>M</i> = 40.7 (<i>SD</i> = 8.56) | <i>M</i> = 24.6 (<i>SD</i> = 8.74) | $t_{(454)} = 17.1$ | <.001 |
| Center for Epidemiologic Studies Depression Scale — short form | <i>M</i> = 28.6 (<i>SD</i> = 8.99) | <i>M</i> = 12.9 (<i>SD</i> = 8.92) | $t_{(454)} = 16.4$ | <.001 |
| Barratt Impulsiveness Scale — short form | <i>M</i> = 35.9 (<i>SD</i> = 7.46) | <i>M</i> = 30.8 (<i>SD</i> = 5.89) | $t_{(454)} = 7.47$ | <.001 |

for measuring uncertainty and concern with regard to perceiving and identifying feelings and hunger signals. Higher scores represent lower interoceptive awareness. Internal consistency was $\alpha = .918$ in the current study.

Center for Epidemiologic Studies Depression Scale — short form

The 15-item short form of the Center for Epidemiologic Studies Depression Scale (Hautzinger, Bailer, Hofmeister, & Keller, 2012; Radloff, 1977) was used for measuring depressive symptoms in the past week. Higher scores represent higher depressiveness. Internal consistency was $\alpha = .944$ in the current study.

Barratt Impulsiveness Scale — short form

The 15-item short form of the Barratt Impulsiveness Scale (Meule, Vögele, & Kübler, 2011; Spinella, 2007) was used for measuring trait impulsivity. Higher scores represent higher impulsivity. Internal consistency was $\alpha = .810$ in the current study.

Data analyses

The groups were compared regarding frequencies of YFAS2.0 diagnoses with a χ^2 test and regarding continuous study variables with independent *t*-tests. Linear regression analyses were used to examine differential relationships between continuous study variables and ‘food addiction’ symptoms as a function of group. Specifically, moderation analyses were conducted with PROCESS for SPSS (Hayes, 2013), where continuous study variables, group (bulimia vs controls), and their interaction predicted the number of YFAS2.0 symptoms. This was performed for each variable separately (BMI, weight suppression, interoceptive awareness, depressive symptoms, and impulsivity; Table 2). Continuous study variables were mean-centred before calculating the product term.

Results

The groups did not differ in age, but the individuals with bulimia had higher BMI and weight suppression, more YFAS2.0 symptoms, lower interoceptive awareness, higher depression scores, and higher impulsivity (Table 1). All but five individuals with bulimia nervosa received a YFAS2.0 diagnosis (Table 1). Of these, 103 (93.6%) were *severe*, 6 (5.46%) were *moderate*, and 1 (0.91%) was *mild*. A significantly lower proportion of controls received a

Table 2 Results from linear regression analyses, in which the number of ‘food addiction’ symptoms was predicted by continuous study variables, group, and their interactions

| Predictors | YFAS2.0 symptoms | | |
|--|------------------|-----------|----------|
| | <i>b</i> | <i>SE</i> | <i>p</i> |
| Body mass index | | | |
| Body mass index | 0.10 | 0.02 | <.001 |
| Group | 6.78 | 0.29 | <.001 |
| Body mass index \times group | -0.11 | 0.04 | .010 |
| Weight suppression | | | |
| Weight suppression | 0.10 | 0.04 | .026 |
| Group | 6.88 | 0.30 | <.001 |
| Weight suppression \times group | -0.17 | 0.08 | .043 |
| Interoceptive awareness | | | |
| Interoceptive awareness | 0.12 | 0.01 | <.001 |
| Group | 5.44 | 0.42 | <.001 |
| Interoceptive awareness \times group | -0.06 | 0.03 | .070 |
| Depression | | | |
| Depression | 0.09 | 0.01 | <.001 |
| Group | 5.74 | 0.42 | <.001 |
| Depression \times group | -0.04 | 0.03 | .226 |
| Impulsivity | | | |
| Impulsivity | 0.08 | 0.02 | <.001 |
| Group | 6.62 | 0.31 | <.001 |
| Impulsivity \times group | -0.05 | 0.04 | .211 |

Notes: Continuous study variables were mean-centred before calculating the product term. Group represents control participants (coded with 0) and individuals with bulimia (coded with 1).

YFAS2.0 diagnosis (Table 1). Of these, 34 (69.4%) were *severe*, 7 (14.3%) were *moderate*, and 8 (16.3%) were *mild*.

Lower interoceptive awareness, higher depression scores, and higher impulsivity were associated with more YFAS2.0 symptoms, independent of group (Table 2). However, there were significant interaction effects between group and BMI and between group and weight suppression when predicting the number of YFAS2.0 symptoms (Table 2). Body mass was positively associated with the number of YFAS2.0 symptoms in the controls ($b = 0.13$, $SE = 0.03$, $p < .001$) but not in the individuals with bulimia ($b = 0.02$, $SE = 0.03$, $p = .575$). Similarly, weight suppression was positively associated with the number of YFAS2.0 symptoms in the controls ($b = 0.14$, $SE = 0.06$, $p = .011$) but not in the individuals

with bulimia ($b = -0.03$, $SE = 0.06$, $p = .663$). Note that these results were not due to the difference in BMI between the groups: We also examined these interaction effects after randomly excluding normal-weight controls so that the groups would not differ in BMI and the interactions would be similarly found.

Discussion

In the current study, almost all participants with bulimia (96%) received a YFAS2.0 diagnosis. This result is in line with studies, in which the previous version of the YFAS was used, which similarly found that prevalence rates of 'food addiction' range between 80% and 100% in individuals with bulimia (Gearhardt et al., 2014; Granero et al., 2014; Meule et al., 2014). Thus, it appears that the changes made in the YFAS2.0 do not affect the association between bulimia and addiction-like eating. In the controls without eating disorders, 14% received a YFAS2.0 diagnosis, which is similar to the prevalence rates found with the English version of the YFAS2.0 (approximately 15%; Gearhardt et al., 2016). Thus, although there is a high overlap between eating disorders (particularly bulimia) and 'food addiction', there is also a substantial subset of individuals without eating disorders who receive a YFAS2.0 diagnosis.

There has been considerable interest in the role of 'food addiction' for weight gain and development of obesity (Meule, 2012). While the present study showed that psychological correlates of 'food addiction' symptomatology (interoceptive awareness, depression, and impulsivity), which have been found previously (Granero et al., 2014; Meule & Gearhardt, 2014a), were similarly found in the individuals with bulimia and the controls, associations with BMI and weight suppression were moderated by group. Specifically, higher BMI and weight suppression were associated with YFAS2.0 symptoms in the controls but not in the participants with bulimia. As has been suggested previously (Meule, 2012), it seems plausible that the positive association between BMI and YFAS2.0 scores is alleviated within the population of individuals with bulimia because of their compensatory weight

control behaviours. Specifically, higher 'food addiction' symptomatology is likely related to more frequent purging in individuals with bulimia, which counteracts weight gain.¹ This, and the fact that those with bulimia already display very high levels of weight suppression, may similarly explain the absent association between higher weight suppression and more YFAS2.0 symptoms in the participants with bulimia, although weight suppression was associated with YFAS2.0 symptoms in controls.

Interpretation of the results is, of course, limited by the use of self-report measures in the current study. For example, it is known that self-reported height is usually overestimated, while self-reported weight is underestimated (Connor Gorber, Tremblay, Moher, & Gorber, 2007). However, there are also studies showing that although these discrepancies exist, data are usually sufficiently accurate (Pursey, Burrows, Stanwell, & Collins, 2014). Although preliminary data regarding agreement of EDDS diagnoses with interview measures have been reported (Sysko et al., 2015), it is likely that the EDDS overestimates the prevalence of eating disorder diagnoses. Nevertheless, as the aims of the current study were to report on associations between the YFAS2.0 and other measures (and not on the prevalence of diagnoses of bulimia), we would argue that these associations are probably not much affected by the absolute numbers of diagnoses.

Notwithstanding these limitations, the current study provides first data on the relationship between bulimia and 'food addiction' as assessed with the YFAS2.0. The results show that, similar to the previous version of the YFAS, there is a very strong overlap between bulimic symptomatology and addiction-like eating. This strong overlap needs to be taken into consideration when reporting and interpreting associations between 'food addiction' and body mass.

Acknowledgement

This work was supported by the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation program (ERC-StG-2014 639445 NewEat).

REFERENCES

- American Psychiatric Association (1994). *Diagnostic and statistical manual of mental disorders* (4th ed.). Washington, DC: American Psychiatric Association.
- American Psychiatric Association (2013). *Diagnostic and statistical manual of mental disorders* (5th ed.). Washington, DC: American Psychiatric Association.
- Benton, D. (2010). The plausibility of sugar addiction and its role in obesity and eating disorders. *Clinical Nutrition*, 29, 288–303.
- Connor Gorber, S., Tremblay, M., Moher, D., & Gorber, B. (2007). A comparison of direct vs. self-report measures for assessing height, weight and body mass index: A systematic review. *Obesity Reviews*, 8, 307–326.
- Garner, D. M. (1991). *Eating Disorder Inventory-2*. Professional manual. Odessa, Florida: Psychological Assessment Resources.
- Gearhardt, A. N., Boswell, R. G., & White, M. A. (2014). The association of "food addiction" with disordered eating and body mass index. *Eating Behaviors*, 15, 427–433.
- Gearhardt, A. N., Corbin, W. R., & Brownell, K. D. (2009). Preliminary validation of the Yale Food Addiction Scale. *Appetite*, 52, 430–436.
- Gearhardt, A. N., Corbin, W. R., & Brownell, K. D. (2016). Development of the Yale Food Addiction Scale Version 2.0. *Psychology of Addictive Behaviors*, 30, 113–121.
- Granero, R., Hilker, I., Aguera, Z., Jimenez-Murcia, S., Sauchelli, S., Islam, M. A., et al. (2014). Food addiction in a Spanish sample of eating disorders: DSM-5 diagnostic subtype differentiation and validation data. *European Eating Disorders Review*, 22, 389–396.
- Hautzinger, M., Bailer, M., Hofmeister, D., & Keller, F. (2012). *Allgemeine Depressionsskala* (2nd ed.). Göttingen: Hogrefe.
- Hayes, A. F. (2013). *Introduction to mediation, moderation, and conditional process analysis*. New York: The Guilford Press.
- Meule, A. (2012). Food addiction and body-mass-index: A non-linear relationship. *Medical Hypotheses*, 79, 508–511.
- Meule, A. (2015). Back by popular demand: A narrative review on the history of food addiction research. *Yale Journal of Biology and Medicine*, 88, 295–302.
- Meule, A. (2016). German version of the Yale Food Addiction Scale 2.0: Associations with impulsivity, trait food craving, binge eating, dieting success, and body mass. *German Medical Science*. 10.3205/16dges080.

¹Note that the frequency of vomiting was indeed negatively (but not significantly) correlated with BMI ($r = -.166$, $p = .076$) and positively (but not significantly) correlated with the number of YFAS2.0 symptoms ($r = .155$, $p = .098$) in the participants with bulimia in the current study.

- Meule, A., & Blechert, J. (2015). German version of the Eating Disorder Diagnostic Scale — DSM-5 version. <https://www.wordpress.com/resources>.
- Meule, A., & Gearhardt, A. N. (2014a). Five years of the Yale Food Addiction Scale: Taking stock and moving forward. *Current Addiction Reports*, *1*, 193–205.
- Meule, A., & Gearhardt, A. N. (2014b). Food addiction in the light of DSM-5. *Nutrients*, *6*, 3653–3671.
- Meule, A., Vögele, C., & Kübler, A. (2011). Psychometric evaluation of the German Barratt Impulsiveness Scale — Short version (BIS-15). *Diagnostica*, *57*, 126–133.
- Meule, A., von Rezori, V., & Blechert, J. (2014). Food addiction and bulimia nervosa. *European Eating Disorders Review*, *22*, 331–337.
- Pursey, K. M., Burrows, T. L., Stanwell, P., & Collins, C. E. (2014). How accurate is web-based self-reported height, weight, and body mass index in young adults? *Journal of Medical Internet Research*, *16*(1), e4.
- Pursey, K. M., Stanwell, P., Gearhardt, A. N., Collins, C. E., & Burrows, T. L. (2014). The prevalence of food addiction as assessed by the Yale Food Addiction Scale: A systematic review. *Nutrients*, *6*(10), 4552–4590.
- Radloff, L. S. (1977). The CES-D scale: A self-report depression scale for research in the general population. *Applied Psychological Measurement*, *1*, 385–401.
- Rogers, P. J., & Smit, H. J. (2000). Food craving and food “addiction”: A critical review of the evidence from a biopsychosocial perspective. *Pharmacology, Biochemistry and Behavior*, *66*, 3–14.
- Schaumberg, K., Anderson, L. M., Reilly, E. E., Gorrell, S., Anderson, D. A., & Earleywine, M. (2016). Considering alternative calculations of weight suppression. *Eating Behaviors*, *20*, 57–63.
- Spinella, M. (2007). Normative data and a short form of the Barratt Impulsiveness Scale. *International Journal of Neuroscience*, *117*, 359–368.
- Stice, E. (2014). Eating Disorder Diagnostic Scale — DSM-5 version. <https://www.ori.org/sticemeasures>.
- Sysko, R., Glasofer, D. R., Hildebrandt, T., Klimek, P., Mitchell, J. E., Berg, K. C., et al. (2015). The eating disorder assessment for DSM-5 (EDA-5): Development and validation of a structured interview for feeding and eating disorders. *International Journal of Eating Disorders*, *48*, 452–463.
- Thiel, A., Jacobi, C., Horstmann, S., Paul, T., Nutzinger, D. O., & Schussler, G. (1997). German translation of the Eating Disorder Inventory EDI-2. *Psychotherapie, Psychosomatik, Medizinische Psychologie*, *47*, 365–376.
- Wilson, G. T. (2010). Eating disorders, obesity and addiction. *European Eating Disorders Review*, *18*, 341–351.
- Ziauddeen, H., & Fletcher, P. C. (2013). Is food addiction a valid and useful concept? *Obesity Reviews*, *14*, 19–28.